

Non-linear Semigroups in Hilbert Spaces used in Petroleum Reservoir Engineering

E.G. Ladopoulos
Interpaper Research Organization
8, Dimaki Str.
Athens, GR - 106 72, Greece
eladopoulos@interpaper.org

Abstract

Non-linear semigroups are used in order to prove the existence and uniqueness of solutions for the non-linear partial differential equation defined in Hilbert spaces and derived from the general porous medium analysis. The above equation is used in well test analysis in petroleum reservoir engineering for the determination of the properties of the reservoir materials. By this method is estimated the size of the oil reserves after their exploration. Beyond the above, the existence and uniqueness of solutions for the non-linear porous medium equation is proved, by presenting some general boundary conditions. Also, some properties of the solutions for the above non-linear partial differential equation are finally proved.

2010 Mathematics Subject Classification : 65L10, 65R20.

Key Word and Phrases

Non-linear Semigroups, Hilbert Spaces, Non-linear Partial Differential Equation, Porous Medium Equation, Petroleum Reservoir Engineering.

1. Introduction

Over the past years an increasing interest was realized on studying non-linear semigroups in general Banach spaces associated with the existence and uniqueness theory of partial differential equations arising in a big level of problems of mathematical physics and engineering. The study of the non-linear semigroups was derived directly from the examination of non-linear parabolic equations and from various non-linear boundary value problems.

The first work on semigroups was published by A.V.Balakrishnan [1], when studying fractional powers of closed operators. Beyond the above, some years later T.Kato [2] studied non-linear semigroups in connection with evolution equations, while Y.Komura [3], [4] studied non-linear semigroups defined in Hilbert spaces.

On the other hand, K.Sato [5] investigated non-negative contraction semigroups in Banach spaces, while a general theory of non-linear semigroups was investigated by M.G. Crandall et al. [6] - [8]. Moreover, J.Watanabe [9], [10] studied semigroups of non-linear operators on closed convex sets and H.Brezis et al. [11], [12] introduced a general semigroups formulation.

At the same time, M.Iannelli [13] studied non-linear semigroups on cones of a non-reflexive Banach space, while J.Mermin [14] and S.Oharu [15] investigated general theories of non-linear semigroups. Beyond the above, I.Miyadera [16] studied semigroups of non-linear operators and B.K.Quinn [17] investigated semigroups in L_1 spaces.

Furthermore, Y.Konishi [18] studied non-linear semigroups associated with some partial differential equations and U.Westphal [19] and S.Aizawa [20] investigated some formulations for non-linear semigroups. On the other hand, T.Kurtz [21] studied semigroups of non-linear operators applied to gas kinetics, while R.Bruck [22] investigated asymptotic convergence of non-linear contraction semigroups in Hilbert spaces.

The theory of non-linear semigroups was generated by Y.Kobayashi [23], [24] and a monograph on the above subject was written by V. Barbu [25]. Moreover, J.M. Ball [26] studied strongly continuous semigroups, while B.C.Burch [27] investigated a semigroup treatment of the Hamilton - Jacobi equations in several space variables.

At the same time, J.H.Lightbourne and R.H.Martin [28] investigated relatively continuous perturbations of analytic semigroups, when A.T.Plant [29] studied non-linear semigroups of translations in Banach spaces generated by functional differential equations.

The theory of non-linear semigroups on general Banach spaces was further investigated by J.B.Baillon [30] and A.Pazy [31], [32], while J.A.Goldstein [33] wrote a monograph on semigroups of linear operators with some general applications. Furthermore, a monograph on non-linear evolution operators and semigroups was written by N.H.Pavel [34].

By the current investigation the non-linear semigroups are used in order to prove the existence and uniqueness of solutions for the non-linear partial differential equation defined in Hilbert spaces. This differential equation is derived from the general theory of porous medium analysis. The porous medium equation was recently used by E.G.Ladopoulos [35] - [37] in well test analysis in the petroleum reservoir engineering for the determination of the properties of the reservoir materials. The above theory together with "Real-time Expert Seismology" was used for the exploration of land and marine oil reserves, as an extension of the non-linear theories investigated by E.G.Ladopoulos et al. during the last two decades. [38] - [49].

Furthermore, in the present report the existence and uniqueness of solutions for the non-linear porous medium equation is investigated, by using a method of non-linear semigroups. Finally, some properties of the solution for the above non-linear differential equation are proved.

2. General Porous Medium Analysis

Theorem 2.1

Suppose that oil flows through a porous medium, that occupies the domain B , which is bounded in R^3 . Denote by $u=u(x,t)$ the density of the oil and by $p=p(x,t)$ its pressure at the point $x = (x_1, x_2, x_3) \in B$ at time t . (Fig.1)




Fig. 1 A bounded domain B in R^3 with small boundary Γ inside which flows a gas with density $u=u(x,t)$ and pressure $p=p(x,t)$ at the point $x = (x_1, x_2, x_3) \in B$ at time t .

Then, the porous medium equation is equal to:

$$\frac{\partial u(x,t)}{\partial t} = \mu \nabla^2 [u(t,x)]^m \quad (2.1)$$

where ∇^2 denotes the Laplace operator:

$$\nabla^2 = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \frac{\partial^2}{\partial x_3^2} \quad (2.2)$$

E.G. Ladopoulos

and μ is equal to:

$$\mu = a\lambda p_0 / \xi(a+1) \quad (2.3)$$

with p_0 a constant, $a \geq 1, \xi > 0$ the viscosity of the medium, $\lambda > 0$ its permeability and $m = a + 1 \geq 2$.

Proof

Let us have:

$$p = p_0 u^a \quad (2.4)$$

where p_0 denotes a constant and $a \geq 1$.

Moreover, if $v = (v, t, u)$ denotes the velocity of the oil, then Darcy's law gives:

$$\xi \mathbf{v} = -\lambda \nabla p \quad (2.5)$$

where ξ denotes the viscosity and λ the permeability of the medium.

Hence, by combining eqs (2.4) and (2.5) we obtain:

$$\xi \mathbf{v} = -\lambda \nabla p_0 u^a \quad (2.6)$$

Furthermore, the dynamic of gas is given by the following conservation law:

$$f \frac{\partial u}{\partial t} + \nabla \cdot (u \mathbf{v}) = 0 \quad (2.7)$$

where f denotes the porosity of the solid, $0 < f < 1$.

Moreover, it is well known that:

$$\nabla \cdot (u \nabla u^a) = \frac{a}{a+1} \nabla^2 u^{a+1} \quad (2.8)$$

Thus, by combining eqs (2.4), (2.5), (2.6), (2.7) and (2.8) we obtain the required formula (2.1).

3. Existence and Uniqueness Theorems for Non-linear Partial Differential Equations in L_2 , by using Non-linear Semigroups

Definition 3.1

Let F a nonempty subset of a Banach space B . Then, a semigroup on F is a function S on R_0^+ such that $S(t) : F \rightarrow F$ for each $t \geq 0$ with the following properties:

$$S(0) = 1 \quad (3.1)$$

$$S(t+s) = S(t)S(s), \quad t, s \geq 0 \quad (3.2)$$

$$\lim_{t \rightarrow 0} S(t)x = x, \quad \forall x \in F \quad (3.3)$$

Theorem 3.1

Consider by B a bounded domain in R^3 , with smooth boundary Γ and $u=u(x,t)$ the temperature function at the point $x = (x_1, x_2, x_3) \in B$ at time t . (Fig.1)

Then, the porous medium equation:

$$\frac{\partial u(x,t)}{\partial t} = \mu \nabla^2 [u(t,x)]^m \quad \text{in }]0, +\infty[, x \in B \quad (3.4)$$

with the boundary conditions:

$$u(t,x) = 0 , \quad \text{on }]0, +\infty[, x \in \Gamma \quad (3.5)$$

$$u(0,x) = u_0(x) , \quad \text{in } B \quad (3.6)$$

where μ is given by (2.3), $m \geq 2$ and $u_0 \in L_2(B)$, has a unique solution $u(t,x) = (S(t)u_0)(x)$, $x \in B$, $t \geq 0$, $u \in C^n \{]0, \infty[; L_2(B)\}$, $n = 1, 2, \dots$

Proof

As it is well known, the Laplace operator ∇^2 is self-adjoint in $L_2(B)$.

Hence, for $u, w \in D(\nabla^2)$ we have:

$$\langle \nabla^2 u, w \rangle = \int_B (\nabla^2 u) w dx \quad (3.7)$$

which is equal to:

$$\int_B \{\nabla^2 u\} w dx = - \int_B (\nabla u)(\nabla w) dx \quad (3.8)$$

and finally to:

$$- \int_B (\nabla u)(\nabla w) dx = \int_B u \nabla^2 w dx = \langle u, \nabla^2 w \rangle \quad (3.9)$$

Thus, from eqs (3.7), (3.8) and (3.9) we obtain:

$$\langle \nabla^2 u, w \rangle = \langle u, \nabla^2 w \rangle \quad (3.10)$$

from which follows that ∇^2 is symmetric and maximal monotone in L_2 and thus $\nabla^2 = \nabla^{2*}$ and $u(t) \in D(\nabla^2)^n$, $n = 0, 1, 2, \dots, t > 0$.

So, the solution $u(t,x) = (S(t)u_0)(x)$ has the property $u \in C^n \{]0, \infty[; D(\nabla^2)^r\}$ for every n , with $r = 0, 1, 2, \dots$ and finally $u \in C^n \{]0, \infty[; C^n(B)\}$, $n = 0, 1, 2, \dots$

Thus, as the solution $u(t,x) = (S(t)u_0)(x)$ tends in $L_2(B)$ to an equilibrium point, then finally follows that: $u \in C^n \{]0, \infty[; L_2(B)\}$, $n = 1, 2, \dots$

Theorem 3.2

Consider the non-linear partial differential porous medium equation (3.4) with the boundary conditions (3.5) and (3.6). Then, the solution $u(t, x) = (S(t)u_0)(x)$ of the above problem, with $u_0 \in L_2(B)$, satisfies the following properties:

$$\frac{1}{2} \|u(T)\|_{L_2}^2 + \int_0^T \|\nabla u(t)\|_{L_2}^2 dt = \frac{1}{2} \|u_0\|_{L_2}^2, \quad \forall T > 0 \quad (3.11)$$

and:
$$\|S(t)u_0\|_{L_p} \leq \|u_0\|_{L_p}, \quad \forall u_0 \in L_p(B), \quad 1 \leq p < \infty \quad (3.12)$$

Proof

Let us consider the following function:

$$g(t) = \frac{1}{2} \|u(t)\|_{L_2}^2 \quad (3.13)$$

Then, we have:

$$g'(t) = \langle u(t), u'(t) \rangle = \int_B u \nabla^2 u dx \quad (3.14)$$

from which follows:

$$\int_B u \nabla^2 u dx = - \int_B |\nabla u(t, x)|^2 dx \quad (3.15)$$

Hence, by integrating over $[\varepsilon, T]$ with $0 < \varepsilon < T$ and letting $\varepsilon \rightarrow 0$, then we obtain (3.11).

Furthermore, by multiplying (3.4) by $|u(t, x)|^{p-2} u(t, x)$ for the case $m = 1$ and taking into account that:

$$\nabla(|u|^{p-2} u) = (p-1)|u|^{p-2} \nabla u \quad (3.16)$$

then, one has:

$$\frac{1}{p} \frac{d}{dt} \|u(t)\|_{L_p}^p = \int_B (\nabla^2 u) u |u|^{p-2} dx \quad (3.17)$$

which finally is equal to:

$$\int_B (\nabla^2 u) u |u|^{p-2} dx = - (p-1) \int_B |\nabla u|^2 |u(t, x)|^{p-2} dx \leq 0 \quad (3.18)$$

from which follows the required (3.12).

4. Conclusions

A general porous medium analysis was presented, by considering an oil which flows through a porous medium which occupies a closed domain, bounded in R^3 . This problem is reduced to the solution of a non-linear partial differential equation, under some general boundary conditions.

Consequently, the existence and uniqueness of solutions was proved, for the above non-linear porous medium equation defined in Hilbert spaces, by using a technique based on non-linear semigroups. This method has clearly simplified the proofs of the existence and uniqueness theorems and has clarified the simpleness to other methods of mathematical analysis.

Finally, some properties for the solutions of the non-linear partial differential equations were proved, when these are defined in Hilbert spaces. These properties were further generalized in L_p spaces, by proving a corresponding inequality for the solution of the non-linear porous medium equation.

References

1. Balakrishnan A.V., 'Fractional powers of closed operators and the semigroups generated by them', *Pacific J. Math.*, **10** (1960), 419-437.
2. Kato T., 'Nonlinear semigroups and evolution equations', *J. Math. Soc. Japan*, **19** (1967), 308-520.
3. Komura Y., 'Nonlinear semigroups in Hilbert space', *J. Math. Soc. Japan*, **19** (1967), 493-507.
4. Komura Y., 'Differentiability of nonlinear semigroups', *J. Math. Soc. Japan*, **21** (1969), 375-402.
5. Sato K., 'On the generators of non-negative contraction semigroups in Banach lattices', *J. Math. Soc. Japan*, **20** (1968), 431-436.
6. Crandall M.G. and Pazy A., 'Semi-groups of nonlinear operators and dissipative sets', *J. Funct. Anal.*, **3** (1969), 376-418.
7. Crandall M.G. and Liggett T. M., 'Generation of semigroups of nonlinear transformations on general Banach spaces', *Amer. J. Math.*, **93** (1971), 265 -298.
8. Crandall M.G., 'A generalized domain for semigroups generators', *Proc. Amer. Math. Soc.*, **37** (1973), 434-440.
9. Watanabe J., 'Semigroups of nonlinear operators on closed convex sets', *Proc. Japan. Acad. Sci.*, **45** (1969), 219 - 223.
10. Watanabe J., 'Autonomous nonlinear functional differential equations and nonlinear semigroups', *J. Math. Anal. Appl.*, **46** (1974), 1-12.
11. Brezis H. and Pazy A., 'Semigroups of nonlinear contractions on convex sets', *J. Funct. Anal.*, **6** (1970), 367-383.
12. Brezis H., 'Operateurs Maximaux Monotones et Semigroups de Contractions dans les Espaces de Hilbert', *Math. Studies*, 5, North Holland, 1973.
13. Ianelli M., 'Non-linear semigroups on cones of a non reflexive Banach space', *Boll. Un. Mat. Ital.*, **3** (1970), 412-419.
14. Mermin J., 'On exponential limit formula and nonlinear semigroups', *Trans. Amer. Math. Soc.*, **150** (1970) 469-476.
15. Oharu S., 'On the generation of semigroups of nonlinear contractions', *J. Math. Soc. Japan*, **22** (1970), 526-550.
16. Miyadera J., 'Some remarks on semi-groups of nonlinear operators', *Tohoku Math. J.*, **23** (1971), 245 - 258.
17. Quinn B. K., 'Solutions with shocks: An example of L_1 contractive semigroups', *Comm. Pure Appl. Math.*, **24** (1971), 125-132.
18. Konishi Y., 'On $u_t = u_{xx} - F(u_x)$ and the differentiability of the nonlinear semigroups associated with it', *Proc. Japan. Acad.*, **48** (1972), 281-286.
19. Westphal U., 'Sur la saturation pour des semigroups non - lineaires', *C.R.Akad. Sci. Paris*, **274** (1972), 1351-1353.
20. Aizawa S., 'A semigroup treatment of the Hamilton - Jacobi equation in one space variable', *Hiros. Math. J.*, **3** (1973), 367-386.
21. Kurtz T., 'Convergence of sequences of semigroups of nonlinear operators with an application to gas kinetics', *Trans. Amer. Math. Soc.*, **186** (1973), 259-272.
22. Bruck R., 'Asymptotic convergence of nonlinear contraction semigroups in Hilbert space', *J. Funct. Anal.*, **18** (1975), 15-26.

E.G. Ladopoulos

23. Kobayashi Y., 'Difference approximation of Cauchy problems for quasi-dissipative operators and generation of nonlinear semigroups', *J. Math. Soc. Japan*, **27** (1975), 641- 663.
24. Kobayashi Y., 'A remark on convergence of nonlinear semigroups', *Proc. Japan Acad., Ser. A*, **55** (1979), 45 - 48.
25. Barbu V., '*Nonlinear Semigroups and Differential Equations in Banach Spaces*', Noordhoff, Leyden, Netherlands, 1976.
26. Ball J. M., 'Strongly continuous semigroups, weak solutions and the variation of constants formula', *Proc. Amer. Math. Soc.*, **63** (1977), 370 -373.
27. Burch B. C., 'A semigroup treatment of the Hamilton - Jacobi equations in several space variables', *J. Diff. Eqns*, **23** (1977), 102-124.
28. Lightbourne J. H. and Martin R. H., 'Relatively continuous nonlinear perturbations of analytic semigroups', *Nonlin. Anal*, **1** (1977), 277-292.
29. Plant A.T., 'Nonlinear semigroups of translations in Banach space generated by functional differential equation', *J. Math. Anal. Appl.*, **60** (1977), 67-74.
30. Baillon J.B., 'Generateurs et semi-groupes dans les espaces de Banach uniformement lisses', *J. Funct. Anal.*, **29** (1978), 199-213.
31. Pazy A., 'The Lyapounov method for semigroups of nonlinear contractions in Banach spaces', *J. Analyse Math.*, **40** (1981), 239-262.
32. Pazy A., '*Semigroups of Linear Operators and Applications to Partial Differential Equations*', Springer, Berlin, 1983.
33. Goldstein J. A., '*Semigroups of Linear Operators and Applications*', Oxford University Press, Oxford, 1985.
34. Pavel M. M., '*Nonlinear Evolution Operators and Semigroups*', Springer, Berlin, 1987.
35. Ladopoulos E.G., 'Non-linear singular integral representation for petroleum reservoir engineering', *Acta Mech.*, **220** (2011), 247-253.
36. Ladopoulos E.G., 'Petroleum reservoir engineering by non-linear singular integral equations', *Mech. Engng Res.*, **1** (2011), 1-10.
37. Ladopoulos E.G., 'Hydrocarbon reserves exploration by Real-time expert seismology and non-linear singular integral equations', *Int. J. Oil Gas Coal Tech.*, **5** (2012), 299-315.
38. Ladopoulos E.G., 'Non-linear singular integral representation for unsteady inviscid flowfields of 2-D airfoils', *Mech. Res. Commun.*, **22** (1995), 25 – 34.
39. Ladopoulos E.G., 'Non-linear singular integral computational analysis for unsteady flow problems', *Renew. Energy*, **6** (1995), 901 – 906.
40. Ladopoulos E.G. and Zisis V.A., 'Non-linear singular integral approximations in Banach spaces', *Nonlin. Anal., Theor. Math. Appl.*, **26** (1996), 1293 – 1299.
41. Ladopoulos E.G. and Zisis V.A., 'Existence and uniqueness for non-linear singular integral equations used in fluid mechanics', *Appl. Math.*, **42** (1997), 345 – 367.
42. Ladopoulos E.G., 'Non-linear singular integral representation analysis for inviscid flowfields of unsteady airfoils', *Int. J. Non-Lin. Mech.*, **32** (1997), 377 – 384.
43. Ladopoulos E.G., 'Collocation approximation methods for non-linear singular integro-differential equations in Banach Spaces', *J. Comp. Appl. Math.*, **79** (1997), 289 – 297.
44. Ladopoulos E.G., 'Non-linear multidimensional singular integral equations in 2-dimensional fluid mechanics analysis', *Int. J. Non-Lin. Mech.*, **35** (2000) , 701-708.
45. Ladopoulos E.G. and Zisis V.A., 'Non-linear finite-part singular integral equations arising in two-dimensional fluid mechanics', *Nonlin. Anal., Th. Meth. Appl.*, **42** (2000), 277-290.
46. Ladopoulos E.G., '*Singular Integral Equations, Linear and Non-Linear Theory and its Applications in Science and Engineering*', Springer Verlag, New York, Berlin, 2000.
47. Ladopoulos E.G., 'Non-linear unsteady flow problems by multidimensional singular integral representation analysis', *Int. J. Math. Math. Scien.*, **2003** (2003), 3203-3216.
48. Ladopoulos E.G., 'Non-linear two-dimensional aerodynamics by multidimensional singular integral computational analysis', *Forsh. Ingen.*, **68** (2003), 105-110.
49. Ladopoulos E.G., 'Unsteady inviscid flowfields of 2-D airfoils by non-linear singular integral computational analysis', *Int. J. Nonlin. Mech.*, **46** (2011), 1022-1026.